

TURAN EXTREMUM PROBLEM FOR PERIODIC FUNCTION WITH SMALL SUPPORT

D.V. GORBACHEV, A.S. MANOSHINA

ABSTRACT. We consider an extremum problem posed by Turan. The aim of this problem is to find a maximum mean value of 1-periodic continuous even function such that sum of Fourier coefficient modules for this function is equal to 1 and support of this function lies in $[-h, h]$, $0 < h \leq 1/2$. We show that this extremum problem for rational $h = p/q$ is equivalent two finite-dimensional linear programming problems. Here there are exact results for rational $h = 2/q$, $h = p/(2p+1)$, $h = 3/q$, and asymptotic equalities.

Suppose $0 < h \leq 1/2$; then by $K(h)$ we denote a class of continuous 1-periodic even functions $f(x)$, where $f(x)$ satisfies following conditions:

- 1) $f(x) = \sum_{n=0}^{\infty} a_n \cos(2\pi nx)$;
- 2) $a_n \geq 0$ for all $n = 0, 1, 2, \dots$;
- 3) $f(0) = \sum_{n=0}^{\infty} a_n = 1$;
- 4) $f(x) = 0$, for $h \leq |x| \leq 1/2$.

For arbitrary h this class $K(h)$ is non-empty. For example, if 1-periodic function $\varphi(x)$ possesses the value $\max\{1 - h^{-1}|x|, 0\}$ on the interval $[-1/2, 1/2]$ and has expansion in Fourier series [1]

$$\varphi_h = h + 2h \sum_{n=1}^{\infty} \left(\frac{\sin(\pi nh)}{\pi nh} \right)^2 \cos(2\pi nx),$$

then this function belongs to $K(h)$.

We estimate the following value

$$A(h) = \sup_{f \in K(h)} a_0 = \sup_{f \in K(h)} \int_{-h}^h f(x) dx.$$

In 1970 P. Turan posed this extremum problem in connection with applications in the theory of number. This problem is connection with van der Corput sets. Other applications in the analytic number theory your can see in [2].

In [1], S.B. Stechkin proved that $A(1/q) = 1/q$ for $h = 1/q$ ($q = 2, 3, \dots$). In this case the extremal function is $\varphi(x)$.

For $h \neq 1/q$ $A(h) > h$ (A.Y. Popov, verbal information).

Theorem 1 ([3]). $A(h) = h + O(h^3)$ as $h \rightarrow 0$. Asymptotically an extremal function is $\varphi(x)$. The constant in O is no more than 16.

Bellow we consider the problem of Turan for arbitrary ration h . Our main results are following.

Date: 11/27/2001.

This research was supported by the RFBR unedr grant no. 00-01-00644, 01-01-06190 and 02-01-06563.

This work was published in Tchebyshev Collection, TPSU (Russia), **2** (2001), 31–40.

Suppose $p, q \in \mathbb{N}$, $(p, q) = 1$ and $2p \leq q$. We consider two problems of linear programming.

$$\begin{aligned} 1. \text{ If } & \sum_{r=0}^{q-1} s_r = 1, \\ & \sum_{r=0}^{q-1} s_r \cos(2\pi rk/q) = 0, \quad k = p, p+1, \dots, q-p, \\ & s_r \geq 0, \quad r = 0, 1, \dots, q-1; \\ & S_1(p, q) = \max_s s_0 \quad \text{is to be found.} \end{aligned}$$

$$\begin{aligned} 2. \text{ If } & 1 + 2 \sum_{k=1}^{p-1} b_k \cos(2\pi rk/q) \geq 0, \quad r = 0, 1, \dots, q-1; \\ & S_2(p, q) = \max_b \frac{1}{q} \left(1 + 2 \sum_{k=1}^{p-1} b_k \right) \quad \text{is to be found.} \end{aligned}$$

Theorem 2. Suppose $p, q \in \mathbb{N}$, $(p, q) = 1$ and $2p \leq q$; then

$$A(p/q) = S_1(p, q) = S_2(p, q).$$

Extremal function is 1-periodic piece linear function $\varphi_{p,q}(x)$ with vertex at points

$$(\pm k/q, b_k^*), \quad k = 0, 1, \dots, p \quad b_0^* = 1, \quad b_p^* = 0.$$

and

$$\varphi_{p,q}(x) = \begin{cases} b_k^* + (b_k^* - b_{k+1}^*)(k - q|x|), & k/q \leq |x| < (k+1)/q, \\ 0, & p/q \leq |x| < 1/2. \end{cases}$$

Here b_k^* ($k = 1, 2, \dots, p-1$) are values of the optimal schedule in linear programming problem 2.

To prove Theorem 2, we need several lemmas.

Lemma 1. $A(p/q) \leq S_1(p, q)$.

Proof. By $f(x)$ denote an arbitrary function of $K(p/q)$. For $f(x)$ following conditions are hold.

1. $f(k/q) = 0$, $k = p, p+1, \dots, q-p$ (from fourth property of $K(p/q)$). For expansion in Fourier series we get

$$\sum_{n=0}^{\infty} a_n \cos(2\pi nk/q) = 0, \quad k = p, p+1, \dots, q-p. \quad (1)$$

Now we have $n = q\nu + r$, where $\nu = 0, 1, 2, \dots$, $r = 0, 1, \dots, q-1$ are quotient and residue of division n/q respectively. Then $\cos(2\pi nk/q) = \cos(2\pi rk/q)$. Using (1), we get

$$\sum_{r=0}^{q-1} s_r \cos(2\pi rk/q), \quad k = p, p+1, \dots, q-p, \quad (2)$$

where $s_r = \sum_{\nu=0}^{\infty} a_{q\nu+r}$. By second property of $K(h)$, it follows that

$$s_r \geq 0, \quad r = 0, 1, \dots, q-1. \quad (3)$$

2. Since $\sum_{n=0}^{\infty} a_n = 1$, we obtain

$$\sum_{r=0}^{q-1} s_r = 1. \quad (4)$$

3. Clearly, $a_0 \leq s_0$.

Since (2),(3),(4) are conditions of linear programming problem 1, we see that $s_0 \leq S_1(p, q)$. Then $A(p/q) \leq S_1(p, q)$.

Lemma 2. $A(p/q) \geq S_2(p, q)$.

Proof. By $K_0(p/q)$ denote subclass of $K(p/q)$ such that it consists of piece linear functions with vertexes at points

$$(\pm k/q, b_k), \quad k = 0, 1, \dots, p, \quad b_0 = 1, \quad b_p = 0.$$

Any such function is given by

$$\varphi(x) = \begin{cases} b_k + (b_k - b_{k+1})(k - q|x|), & k/q \leq |x| < (k+1)/q, \\ 0, & p/q \leq |x| < 1/2. \end{cases} \quad (5)$$

Its expansion in Fourier series is

$$\varphi(x) = \sum_{n=0}^{\infty} \alpha_n \cos(2\pi n x),$$

where

$$\alpha_0 = \frac{1}{q} \left(1 + 2 \sum_{k=1}^{p-1} b_k \right), \quad \alpha_n = 2q \left(\frac{\sin(\pi n/q)}{\pi n} \right)^2 \left(1 + 2 \sum_{k=1}^{p-1} b_k \cos(2\pi n k/q) \right).$$

For $\varphi(x) \in K_0(p/q)$ it is necessary to have $\alpha_n \geq 0$, ($n = 0, 1, 2, \dots$), i.e.

$$1 + 2 \sum_{k=1}^{p-1} b_k \cos(2\pi n k/q) \geq 0, \quad n = 0, 1, 2, \dots$$

or

$$1 + 2 \sum_{k=1}^{p-1} b_k \cos(2\pi r k/q) \geq 0, \quad r = 0, 1, \dots, q-1.$$

The subclass $K_0(p/q)$ is non-empty. In particular $\varphi_{p/q}(x) \in K_0(p/q)$. We shall seek function from $K_0(p/q)$ such that its coefficient α_0 is highest possible. For that we solve

$$\alpha_0 = \frac{1}{q} \left(1 + 2 \sum_{k=1}^{p-1} b_k \right) \rightarrow \max$$

under conditions

$$1 + 2 \sum_{k=1}^{p-1} b_k \cos(2\pi r k/q) \geq 0, \quad r = 0, 1, \dots, q-1.$$

This problem is the linear programming problem 2.

Let b_k^* ($k=1, 2, \dots, p-1$) be an optimal schedule in the problem 2. We define function $\varphi_{p,q}(x)$ from $K_0(p/q) \subset K(p/q)$ by formula (5) with coefficients b_k^* . Its mean value is $S_2(p, q)$. Since $\varphi_{p,q} \in K(p/q)$ we have $A(p/q) \geq S_2(p, q)$.

Lemma 3. $S_1(p, q) = S_2(p, q)$.

Proof. Let us consider trigonometric sums from problem 2

$$b_k = \sum_{r=0}^{q-1} s_r \cos(2\pi r k/q), \quad k = 0, 1, \dots, q-1,$$

where

$$b_0 = 1, \quad b_p = b_{p+1} = \dots = b_{q-p} = 0. \quad (6)$$

Using discrete cosine Fourier inversion, we get

$$\frac{s_r + s_{q-r}}{2} = \frac{1}{q} \sum_{k=0}^{q-1} b_k \cos(2\pi rk/q), \quad r = 0, 1, \dots, q-1, \quad s_0 = s_q.$$

Since $\cos(2\pi r(q-k)/q) = \cos(2\pi rk/q)$, by (6) it follows that

$$\frac{s_r + s_{q-r}}{2} = \frac{1}{q} \left(1 + 2 \sum_{k=1}^{p-1} b_k \cos(2\pi rk/q) \right).$$

Therefore the finding value $S_1(p, q)$ is equivalent following problem:

$$\max s_0 = \max_b \frac{1}{q} \left(1 + 2 \sum_{k=1}^{p-1} b_k \right)$$

under conditions

$$q \frac{s_r + s_{q-r}}{2} = 1 + 2 \sum_{k=1}^{p-1} b_k \cos(2\pi rk/q) \geq 0, \quad r = 0, 1, \dots, q-1.$$

But this is the linear programming problem 2 and lemma follows.

Thus from lemmas 1–3, we get Theorem 2. Using this theorem, we can find value $A(p/q)$ for fast values p and q .

Theorem 3. For $p = 2$ and $h = 2/q$, $q = 5, 7, \dots$

$$A(2/q) = \frac{1 + \cos(\pi/q)}{q \cos(\pi/q)}.$$

Proof. By Theorem 2, so that $A(2/q) = S_2(2, q)$. We have

$$A(2/q) = \frac{1}{q} \max_b (1 + 2b), \quad 1 + 2b \cos(2\pi r/q) \geq 0, \quad r = 0, 1, \dots, q-1.$$

For $r = 0, 1, \dots, q-1$ and odd q we have $\min(\cos(2\pi r/q)) = -\cos(\pi/q)$. Therefore

$$\max 2b = 2b_1^* = \frac{1}{\cos(\pi/q)} \tag{7}$$

and

$$A(2/q) = \frac{1}{q} \left(1 + \frac{1}{\cos(\pi/q)} \right) = \frac{1 + \cos(\pi/q)}{q \cos(\pi/q)}.$$

An extremal function is the piece linear function $\varphi_{2,q}(x)$ (5) with vertex at points $(0, 1)$, $(\pm 1/q, b_1^*)$, $(\pm 2/q, 0)$ (7).

Theorem 4. For $h = \frac{p}{2p+1}$, $p = 1, 2, \dots$

$$A\left(\frac{p}{2p+1}\right) = \frac{\cos(\frac{\pi}{2p+1})}{1 + \cos(\frac{\pi}{2p+1})}.$$

Proof. From Theorem 2, we get $A(\frac{p}{2p+1}) = S_1(p, 2p+1)$ and

$$A\left(\frac{p}{2p+1}\right) = \max_s s_0$$

under conditions

$$\sum_{r=0}^{q-1} s_r = 1, \quad (8)$$

$$\sum_{r=0}^{q-1} s_r \cos(2\pi rk/q) = 0, \quad k = p, p+1, \quad (9)$$

$$s_r \geq 0, \quad r = 0, 1, \dots, q-1. \quad (10)$$

From (8) we have $s_1 = 1 - s_0 - \sum_{r=2}^{q-1} s_r$. For $k = p$ and (9)

$$s_0 + \left(1 - s_0 - \sum_{r=2}^{q-1} s_r\right) \cos(2\pi p/q) + \sum_{r=2}^{q-1} s_r \cos(2\pi rp/q) = 0.$$

Therefore

$$s_0 (1 - \cos(2\pi p/q)) = -\cos(2\pi p/q) + \sum_{r=2}^{q-1} s_r (\cos(2\pi p/q) - \cos(2\pi rp/q)). \quad (11)$$

Since for $q = 2p+1$

$$\cos\left(\frac{2\pi p}{q}\right) = -\cos\left(\frac{\pi}{2p+1}\right)$$

and

$$\cos\left(\frac{2\pi p}{q}\right) - \cos\left(\frac{2\pi rp}{q}\right) = -\cos\left(\frac{\pi}{2p+1}\right) - (-1)^r \cos\left(\frac{\pi r}{2p+1}\right) \leq 0, \\ r = 2, 3, \dots, q-1,$$

from (10)–(11) we get

$$s_0 \left(1 + \cos\left(\frac{\pi}{2p+1}\right)\right) = \cos\left(\frac{\pi}{2p+1}\right) + \\ + \sum_{r=2}^{q-1} s_r \left(-\cos\left(\frac{\pi}{2p+1}\right) - (-1)^r \cos\left(\frac{\pi r}{2p+1}\right)\right) \leq \cos\left(\frac{\pi}{2p+1}\right) \quad (12)$$

Therefore

$$\max_s s_0 \leq \frac{\cos\left(\frac{\pi}{2p+1}\right)}{1 + \cos\left(\frac{\pi}{2p+1}\right)}.$$

In (12) we have equality for

$$s_0^* = \frac{\cos\left(\frac{\pi}{2p+1}\right)}{1 + \cos\left(\frac{\pi}{2p+1}\right)} > 0, \\ s_1^* = \frac{1}{1 + \cos\left(\frac{\pi}{2p+1}\right)} > 0, \\ s_r^* = 0, \quad r = 2, 3, \dots, q-1.$$

Thus

$$A\left(\frac{p}{2p+1}\right) = \frac{\cos\left(\frac{\pi}{2p+1}\right)}{1 + \cos\left(\frac{\pi}{2p+1}\right)}$$

and extremal function is piece linear function $\varphi_{p,2p+1}(x)$ with vertex at points $(\pm k/q, b_k^*)$, $k = 0, 1, \dots, p$, where (see proof of lemma 3)

$$b_k^* = \frac{\cos\left(\frac{\pi}{2p+1}\right) + \cos\left(\frac{2\pi k}{2p+1}\right)}{1 + \cos\left(\frac{\pi}{2p+1}\right)}.$$

Notice that asymptotically (as $p \rightarrow \infty$) an extremal function is $\cos^2(\pi x)$.

Theorem 5. For $p = 3$ and $h = 3/q$, $q = 7, 8, 10, 11, \dots$

$$A(3/q) = \frac{1}{q} \left(1 + \frac{1 - 2(\cos(2\pi r_0/q) + \cos(2\pi(r_0 + 1)/q))}{1 + 2\cos(2\pi r_0/q)\cos(2\pi(r_0 + 1)/q)} \right),$$

where $r_0 = [q/3]$ is an integer part of $q/3$.

Proof. From Theorem 2, we get

$$A(3/q) = S_2(3, q) = \frac{1}{q} \max_{\alpha} \alpha(0), \quad \alpha(r) \geq 0, \quad r = 0, 1, \dots, q-1, \quad (13)$$

where $\alpha(r)$ is trigonometric polynomial

$$\alpha(r) = 1 + 2b_1 \cos(2\pi r/q) + 2b_2 \cos(4\pi r/q). \quad (14)$$

We define a quadrature formula

$$1 = \frac{1}{q} \sum_{r=0}^{q-1} \alpha(r) = \gamma_0 \alpha(0) + \gamma_1 \alpha(r_0) + \gamma_2 \alpha(r_0 + 1), \quad r_0 = [q/3]. \quad (15)$$

For equality (15) for any trigonometric polynomial (14), it is sufficient to have (15) for basis functions $1, \cos(2\pi r/q), \cos(4\pi r/q)$:

$$\begin{aligned} 1 &= \frac{1}{q} \sum_{r=0}^{q-1} 1 = \gamma_0 + \gamma_1 + \gamma_2, \\ 0 &= \frac{1}{q} \sum_{r=0}^{q-1} \cos(2\pi r/q) = \gamma_0 + \gamma_1 \cos(2\pi r_0/q) + \gamma_2 \cos(2\pi(r_0 + 1)/q), \\ 0 &= \frac{1}{q} \sum_{r=0}^{q-1} \cos(4\pi r/q) = \gamma_0 + \gamma_1 \cos(4\pi r_0/q) + \gamma_2 \cos(4\pi(r_0 + 1)/q). \end{aligned}$$

Since $\gamma_0 \neq 0$, we get combined equations

$$\begin{cases} \gamma_1' \cos(2\pi r_0/q) + \gamma_2' \cos(2\pi(r_0 + 1)/q) = -1, \\ \gamma_1' \cos(4\pi r_0/q) + \gamma_2' \cos(4\pi(r_0 + 1)/q) = -1, \end{cases}$$

where $\gamma_1' = \gamma_1/\gamma_0$ and $\gamma_2' = \gamma_2/\gamma_0$.

We have

$$\gamma_i' = \frac{\Delta_i}{\Delta}, \quad i = 1, 2. \quad (16)$$

Here

$$\begin{aligned} \Delta &= \cos(2\pi r_0/q) \cos(4\pi(r_0 + 1)/q) - \cos(2\pi(r_0 + 1)/q) \cos(4\pi r_0/q) = \\ &= (\cos(2\pi(r_0 + 1)/q) - \cos(2\pi r_0/q))(1 + 2\cos(2\pi r_0/q)\cos(2\pi(r_0 + 1)/q)) < 0 \end{aligned}$$

(from inequalities $q/3 - 1 < r_0 < r_0 + 1 < q/3 + 1$),

$$\begin{aligned} \Delta_1 &= \cos(2\pi(r_0 + 1)/q) - \cos(4\pi(r_0 + 1)/q) = \\ &= 2\sin(\pi(r_0 + 1)/q)\sin(3\pi(r_0 + 1)/q) < 0 \end{aligned}$$

(from inequalities $q/3 < r_0 + 1 < q/3 + 1$),

$$\begin{aligned}\Delta_2 &= \cos(4\pi r_0/q) - \cos(2\pi r_0/q) = \\ &= -2 \sin(\pi r_0/q) \sin(3\pi r_0/q) < 0\end{aligned}$$

(from inequalities $q/3 - 1 < r_0 < q/3$).

Thus, $\gamma'_1, \gamma'_2 > 0$. Therefore all weights γ_i ($i = 0, 1, 2$) have like signs. From $\gamma_0 + \gamma_1 + \gamma_2 = 1$ we get $\gamma_i > 0$ and $\gamma'_1 + \gamma'_2 = (1 - \gamma_0)/\gamma_0$. From (16) we have

$$\frac{1}{\gamma_0} = 1 + \gamma'_1 + \gamma'_2 = 1 + \frac{1 - 2(\cos(2\pi r_0/q) + \cos(2\pi(r_0 + 1)/q))}{1 + 2 \cos(2\pi r_0/q) \cos(2\pi(r_0 + 1)/q)}. \quad (17)$$

Using the quadrature formula (15) for arbitrary trigonometric polynomial (14) ($\alpha(r) \geq 0$, $r = 0, 1, \dots, q - 1$), we get

$$1 = \frac{1}{q} \sum_{r=0}^{q-1} \alpha(r) = \gamma_0 \alpha(0) + \gamma_1 \alpha(r_0) + \gamma_2 \alpha(r_0 + 1).$$

Since $\gamma_1 \alpha(r_0) \geq 0$ and $\gamma_2 \alpha(r_0 + 1) \geq 0$, it follows that $\gamma_0 \alpha(0) \leq 1$ and

$$\alpha(0) \leq \frac{1}{\gamma_0}.$$

The equality reaches for $\alpha^*(r) = 1 + 2b_1^* \cos(2\pi r/q) + 2b_2^* \cos(4\pi r/q)$, where

$$\begin{aligned}b_1^* &= -\frac{\cos(2\pi r_0/q) + \cos(2\pi(r_0 + 1)/q)}{1 + 2 \cos(2\pi r_0/q) \cos(2\pi(r_0 + 1)/q)} > 0, \\ b_2^* &= \frac{1/2}{1 + 2 \cos(2\pi r_0/q) \cos(2\pi(r_0 + 1)/q)} > 0.\end{aligned} \quad (18)$$

An extremal function is the piece linear function $\varphi_{3,q}(x)$ (5) with vertex at points $(0, 1)$, $(\pm 1/q, b_1^*)$, $(\pm 2/q, b_2^*)$, $(\pm 3/q, 0)$ (18). Theorem is proved.

Notice that the main idea of using quadrature formula (15) belong to V.I. Ivanov. In [4] there is an other proof.

Finally, we obtain asymptotic equalities.

Corollary 1. For $h = 2/q \rightarrow 0$, $q = 2\nu + 1$, $\nu = 2, 3, \dots$

$$A(h) = h + \frac{\pi^2}{16} h^3 + \frac{5\pi^4}{768} h^5 + O(h^7).$$

For $h = 3/q \rightarrow 0$, $q = 3\nu + 1$, $\nu = 2, 3, \dots$

$$A(h) = h + \frac{16\pi^2}{243} h^3 - \frac{16\pi^3\sqrt{3}}{2187} h^4 + \frac{448\pi^4}{59049} h^5 + O(h^6),$$

for $h = 3/q \rightarrow 0$, $q = 3\nu + 2$, $\nu = 2, 3, \dots$

$$A(h) = h + \frac{16\pi^2}{243} h^3 + \frac{16\pi^3\sqrt{3}}{2187} h^4 + \frac{448\pi^4}{59049} h^5 + O(h^6).$$

Therefore in Theorem 1 we can not replace O-large by o-small. We express one's thanks to V.I. Ivanov for useful discussions.

REFERENCES

- [1] S.B. Stechkin, *An extremum problem for trigonometric series with non-negative coefficients*, Selected Works: Mathematics [in Russian], Nauka, Moscow, 1998, 244–245.
- [2] S. Konyagin, I. Shparlinski, *Character sums with exponential functions and their applications*, Cambridge Univ. Press, 1999.
- [3] D.V. Gorbachev, *Extremum problem for periodic function supported in ball*, Mathematical Notes, **69** (2001), 346–352.
- [4] A.S. Manoshina, *Extremum problem of Turan for functions with small support*, Izvestiya of the Tula State University. Series Mathematics. Mechanics. Informatics, **6**, N 3 (2000), 113–116.

E-mail address: `dvg@uic.tula.ru`; `manoshin@tula.net`

URL: `http://home.uic.tula.ru/~gd030473`